

GRXB E.Coli

Description:GRXB produced in E.Coli is a single, non-glycosylated polypeptide chain containing 235 amino acids (1-215 a.a.) and having a molecular mass of 26.5kDa.GRXB is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-137

Synonyms:Glutaredoxin-2, Grx2, grxB, b1064, JW1051.

For research use only.

Source:Escherichia Coli.

Physical Appearance:Sterile filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MKLYIYDHCP YCLKARMIFG
LKNIPVELHV LLNDDAETPT RMVGQKQVPI LQKDDSRYP ESMDIVHYVD KLDGKPLLTG
KRSPAIEEWL RKNVGYANKL LLPRFAKSAF DEFSTPAARK YFVDKKEASA GNFADLLAHS
DGLIKNISDD LRALDKLIVK PNAVNGELSE DDIQLFPLLR NLTLVAGINW PSRVADYRDN
MAKQTQINLL SS

Purity:Greater than 95% as determined by SDS-PAGE.

Formulation:

GRXB protein solution (1mg/ml) 20mM Tris-HCl buffer (pH 8.0), 1mM DTT, 10% glycerol and 50mM NaCl.

Stability:

GRXB E.Coli Recombinant although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Glutaredoxin-2 (GrxB) is a member of the glutaredoxin family. Glutaredoxins are small redox enzymes of approximately 100 amino-acid residues which use glutathione as a cofactor. Glutaredoxins are oxidized by substrates, and reduced non-enzymatically by glutathione. GrxB is involved in reducing some disulfides in a coupled system with glutathione reductase. GrxB does not act as hydrogen donor for ribonucleotide reductase.

To place an order, please [Click HERE](#).