www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

PPP1CA Human

Description: PPP1CA Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 350 amino acids (1-330 a.a) and having a molecular mass of 39.7kDa.PPP1CA is fused to a 20 amino acid His-tag at N-terminus & Durified by proprietary chromatographic techniques.

Catalog #:ENPS-172

For research use only.

Synonyms: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit, PP-1A, PPP1CA, PPP1A, PP1alpha.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MSDSEKLNLD SIIGRLLEVQ GSRPGKNVQL TENEIRGLCL KSREIFLSQP ILLELEAPLK ICGDIHGQYY DLLRLFEYGG FPPESNYLFL GDYVDRGKQS LETICLLLAY KIKYPENFFL LRGNHECASI NRIYGFYDEC KRRYNIKLWK TFTDCFNCLP IAAIVDEKIF CCHGGLSPDL QSMEQIRRIM RPTDVPDQGL LCDLLWSDPD KD

Purity: Greater than 80.0% as determined by SDS-PAGE.

Formulation:

PPP1CA protein solution (0.25mg/ml) containing 50mM Tris-HCl buffer (pH8.0), 40% glycerol, 0.15M NaCl, 1mM DTT, 0.1mM PMSF and 1mM MnCl2.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Serine/Threonine-protein phosphatase PP1-alpha catalytic subunit (PPP1CA) is crucial for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. The PPP1CA protein may have a significant role in dephosphorylating substrates such as the postsynaptic density-associated Ca2+/calmodulin dependent protein kinase II.

Biological Activity:

Specific activity: >3,000 units/mg. Enzymatic activity was confirmed by measuring the amount of enzyme hydrolyzing 1 nmole of p-nitrophenyl phosphate (pNPP) per minute at 37C, pH7.5 using 10mM of substrate.

To place an order, please Click HERE.





