www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## IMPDH2 Human

Description: IMPDH2 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 534 amino acids (1-514) and having a molecular mass of 58kDa.IMPDH2 is fused to a 20 amino acid His-tag at N-terminus & Dyrified by proprietary chromatographic techniques.

Catalog #:ENPS-194

For research use only.

Synonyms: Inosine-5'-monophosphate dehydrogenase 2, IMP dehydrogenase 2, IMPD 2, IMPDH 2, IMPDH-II, IMPDH2, IMPD2.

Source: E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MADYLISGGT SYVPDDGLTA QQLFNCGDGL TYNDFLILPG YIDFTADQVD LTSALTKKIT LKTPLVSSPM DTVTEAGMAI AMALTGGIGF IHHNCTPEFQ ANEVRKVKKY EQGFITDPVV LSPKDRVRDV FEAKARHGFC GIPITDTGRM GSRLVGIISS RDIDFLKEEE HDCFLEEIMT KREDLVVAPA GITLKEANEI LQRSKKGKLP IV

Purity: Greater than 90% as determined by SDS-PAGE.

#### Formulation:

The IMPDH2 solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 2mM DTT, 20% glycerol and 150mM NaCl.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

# Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Introduction:

IMPDH2 is a member of the IMPDH/GMPR family. IMPDH2 catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is afterward converted into guanosine-5'-monophosphate. IMPDH2 is the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. IMPDH2 is consequently involved in maintaining cellular guanine deoxy- and ribonucleotide pools required for DNA and RNA synthesis. In addition, IMPDH1 and IMPDH2 are targets for the important immunosuppressive drug, mycophenolic acid (MPA). Furthermore, the IMPDH2 gene is up-regulated in some neoplasms, suggesting it may have a role in malignant transformation.

To place an order, please Click HERE.





