

ACAD8 Human

Description: Acyl-Coenzyme A Dehydrogenase 8 Human Recombinant produced in E.Coli is a non-glycosylated, polypeptide chain containing amino acids 1-415 and having a total molecular mass of 47.7 kda. ACAD8 contains T7 tag at the N-terminus. ACAD-8 is purified by proprietary chromatographic techniques.

Catalog #:ENPS-301

For research use only.

Synonyms: Acyl-CoA dehydrogenase family member 8 mitochondrial, ACAD-8, Isobutyryl-CoA dehydrogenase, Activator-recruited cofactor 42 kDa component, ARC42, FLJ22590.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

Acyl-Coenzyme A Dehydrogenase 8 at a concentration of 0.1mg/ml in 10mM Tris, pH 8.0, 0.1% Triton X-100, 0.002% NaN₃, 10mM DTT.

Stability:

ACAD8 although stable at 4°C for 1 week, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Acyl CoA dehydrogenase is the enzyme used to catalyze the first step of β -oxidation in Fatty acid metabolism. Acyl-coenzyme A (CoA) dehydrogenases (ACADs) are a family of mitochondrial enzymes that catalyze the first dehydrogenation step in the β -oxidation of fatty acyl-CoA derivatives. Several human ACADs exist and all ACADs catalyze the same initial dehydrogenation of the substrate at the beta-carbon atom and require electron transfer flavoprotein as an electron acceptor. The predicted 415-amino acid ACAD8 protein contains many of the residues conserved in most other ACADs, including an active site glutamic acid residue and residues important for tetramer formation.

To place an order, please [Click HERE](#).