

## SlyD E.Coli

**Description:** SlyD Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 196 amino acids and having a molecular mass of 21 kDa.

**Catalog #:** ENPS-345

**Synonyms:** FKBP-Type Peptidyl-Prolyl Cis-Trans Isomerase, SlyD.

For research use only.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered colorless solution.

**Amino Acid Sequence:** MKVAKDLVVS LAYQVRTEDG VLVDESPVSA PLDYLHGHGS  
LISGLETALE GHEVGDKFDV AVGANDAYGQ YDENLVQRVP KDVFIMGVDEL QVGMRFLEET  
DQGPVPEIT AVEDDHVVVD GNHMLAGQNL KFNVEVVAIR EATEEELAAG HVHGAHDHGH  
DHDHDGCCGG HGHDHGHEHG GEGCCGGKGN GGCGCH.

**Purity:** Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

SlyD protein solution contains 20mM Tris pH-7.5.

**Stability:**

SlyD although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Introduction:**

SlyD accession#: NP\_755987 is a putative folding helper protein from the Escherichia coli cytosol, which has N-terminal prolyl isomerase domain of the FKBP type and a most likely unstructured C-terminal tail. SlyD is an important factor in the biosynthesis of the metal cluster in the [NiFe]-hydrogenase enzymes, and exhibits several activities including that of a peptidyl-prolyl isomerase.

**Biological Activity:**

Specific activity is > 220 nmoles/min/mg, and is defined as the amount of enzyme that cleaves 1umole of suc-AAFP-pNA per minute at 25C in Tris-Hcl pH8.0 using chymotrypsin.

**To place an order, please [Click HERE](#).**