www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

UBE2B Human

Description: Ubiquitin Conjugating Enzyme E2B Human Recombinant produced in E.coli is a 19 kDa protein containing 166 amino acids. The UE2B protein contains 6xHis tag and is purified by proprietary chromatographic techniques.

Catalog #:ENPS-347

For research use only.

Synonyms: Ubiquitin-conjugating enzyme E2 B, EC 6.3.2.19, Ubiquitin-protein ligase B, Ubiquitin carrier protein B, HR6B, hHR6B, E2-17 kDa UBC2, HHR6B, RAD6B, E2-17kDa, UBE2B.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered white lyophilized powder.

Amino Acid Sequence:

MHHHHHHAMGOLRSMSTPARRRLMRDFKRLQEDPPVGVSGAPSENNIMOWNAVIFGPEGTPFE DGTFKLVIEFSEEYPNKPPTVRFLSKMFHPNVYADGSICLDILQNRWSPTYDVSSILTSIQSLLDEP NPNSPANSQAAQLYQENKREYEKRVSAIVEQSWNDS.

Purity: Greater than 95.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a 0.2m filtered concentrated (1 mg/ml) solution in 1X PBS and 1mM DTT, pH 7.5.

Stability:

Lyophilized UBE2B although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution UBE2B should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized UBE2B in sterile water not less than 100

Introduction:

This E2 enzyme encodes for the human homolog of the yeast DNA repair gene RAD6, which is induced by DNA damaging agents. UBE2B can conjugate ubiquitin to histone H2A in an E3independent manner in vitro, and is essential for the multi-ubiquitination and degradation of N-end rule substrates. Additionally, UBE2B may have a role in sepsis-induced muscle protein proteolysis and cancer-induced cachexia.

To place an order, please Click HERE.





