www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

NMNAT1 Human

Description: NMNAT1 Recombinant Human produced in E.Coli is a single, non-glycosylated polypeptide chain containing 315 amino acids (1-279 a.a.) and having a molecular mass of 36 kDa. The NMNAT1 is fused to 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

Catalog #:ENPS-391

For research use only.

Synonyms: NMNAT, NMNAT1, PNAT1, Nicotinamide mononucleotide adenylyltransferase 1, NMN adenylyltransferase 1, Nicotinate-nucleotide adenylyltransferase 1, NaMN adenylyltransferase 1, EC=2.7.7.1, EC=2.7.7.18.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSMENS EKTEVVLLAC GSFNPITNMH LRLFELAKDY MNGTGRYTVV KGIISPVGDA YKKKGLIPAY HRVIMAELAT KNSKWVEVDT WESLQKEWKE TLKVLRHHQE KLEASDCDHQ QNSPTLERPG RKRKWTETQD SSQKKSLEPK TKAVPKVKLL CGADLLESFA VPNLWKSEDI TQIVANYGLI CVTRAGNDAQ KF

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

NMNAT1 Human solution containing 20mM Tris pH-8, 0.1M NaCl, 1mM DTT, 1mM EDTA & DTA 20% glycerol.

Stability:

NMNAT1 Human although stable at 4°C for 1 week, should be stored desiccated below -18°C. Please prevent freeze thaw cycles.

Usage:

NeoBiolab's products are furnished forLABORATORY RESEARCHUSEONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

NMNAT1 enzyme is vital for NAD biosynthesis, catalyzing the condensation of nicotinamide mononucleotide (NMN) or nicotinic acid mononucleotide (NaMN) with the AMP moiety of ATP to form NAD or NaAD. NMNAT1 is widely expressed with high levels in skeletal muscle, heart, liver and kidney. This protein appears to have the ability to protect against axonal degeneration following mechanical or toxic insults.

To place an order, please Click HERE.





