www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

GSTT1 Human

Description:GSTT1 Human Recombinant fused with 37 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 277 amino acids (1-240 a.a.) and having a molecular mass of 31.5kDa.The GSTT1 is purified by proprietary chromatographic techniques.

Catalog #:ENPS-436

For research use only.

Synonyms: Glutathione S-transferase theta-1, GST class-theta-1, Glutathione transferase T1-1, GSTT1.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence:MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSHMGL ELYLDLLSQP CRAVYIFAKK NDIPFELRIV DLIKGQHLSD ACAQVNPLKK VPALKDGDFT LTESVAILLY LTRKYKVPDY WYPQDLQARA RVDEYLAWQH TTLRRSCLRA LWHKVMFPVF LGEPVSPQTL AATLAELDVT LQLLEDKFLQ NKAFLTGPHI SLADLVAITE LMHPVGAGCQ VFEGRPKLAT WR

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:
The GSTT1 solution contains 20mM Tris-HCl buffer (pH 8.0) and 10% glycerol.

Stability:
Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drµgs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:
GSTT1 belongs to a superfamily of proteins which catalyze the conjµgation of reduced glutathione to a variety of electrophilic and hydrophobic compounds. GSTT1 is one of the GSTs four main classes: alpha, mu, pi and theta (which includes GSTT1 and GSTT2). GSTT1 is involved in activation and detoxification reactions and catalyzes the conjµgation of industrial chemicals, such as epoxybutane, ethylene oxides, halomethane with glutathione. GSTT1 is found in erythrocytes, at low levels in the liver as well as in Clara and ciliated cells at the alveolar/bronchiolar junction in the lung.The GSTT1 gene is deficient in 38% of the population. The GSTTI enzyme deficiency might influence the individual risk for development of acquired aplastic anemia and acute myeloid leukemia. The presence or absence of the GSTT1 gene is concurrent with GSST1+ (the conjµgator) and GSTT1- (the non-conjµgator) phenotypes correspondingly. The GSTT1+ phenotype is able to catalyze the glutathione conjµgation of dichloromethane. GSTT1-null genotypes are seen as having a higher risk of developing leukoplakia. Germline genetic polymorphism in GSTT1 is linked to breast cancer.

To place an order, please Click HERE.

© 2015 NeoScientific

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)
Request a CATALOG CLICK HERE