www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

Esterase D Human

Catalog #:ENPS-454

For research use only.

Description: Esterase-D Human Recombinant fused with 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 302 amino acids (1-282 a.a.) and having a molecular mass of 33.6kDa.The Esterase-D is purified by proprietary chromatographic techniques.

Synonyms:S-formylglutathione hydrolase, FGH, Esterase D, ESD, FLJ11763, esterase D/formylglutathione hydrolase.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MALKQISSNK CFGGLQKVFE HDSVELNCKM KFAVYLPPKA ETGKCPALYW LSGLTCTEQN FISKSGYHQS ASEHGLVVIA PDTSPRGCNI KGEDESWDFG TGAGFYVDAT EDPWKTNYRM YSYVTEELPQ LINANFPVDP QRMSIFGHSM GGHGALICAL KNPGKYKSVS AFAPICNPVL CPWGKKAFSG YLGTDQSKWK AYDATHLVKS YP

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:
The Esterase-D solution contains 20mM Tris-HCl buffer (pH8.0) and 10% glycerol.

Stability:
Esterase-D althoμgh stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drμgs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:
Formylglutathione hydrolase (Esterase D) is a member of the esterase D family. Esterase D is a serine hydrolase involved in the detoxification of formaldehyde. Esterase D is active toward various substrates including O-acetylated sialic acids, and it may possibly be involved in the recycling of sialic acids. Esterase D is used as a genetic marker for retinoblastoma and Wilson's disease.

To place an order, please Click HERE.

© 2015 NeoScientific