

ALAD Human

Description:ALAD Human Recombinant produced in E. coli is a single polypeptide chain containing 354 amino acids (1-330) and having a molecular mass of 38.8kDa.ALAD is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:ENPS-593

Synonyms:Aminolevulinate delta-dehydratase, ALADH, PBGS, Porphobilinogen synthase, delta-aminolevulinic acid dehydratase, EC 4.2.1.24.

For research use only.

Source:E.coli.

Physical Appearance:Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MGSHMQPQSV LHSYGFHPLL
RAWQTATTTT NASNLIYPIF VTDVPDDIQP ITSLPGVARY GVKRLEEMLR PLVEEGLRCV
LIFGVPSRVP KDERGSAADS EESPAIEAIH LLRKTFPNLL VACDVCLCPY TSHGHCGLLS
ENGAFAEES RQRLAEVALA YAKAGCQVVA PSDMMDGRVE AIKEALMAHG LGNRVSVMSY
SAKFASCFYG PF

Purity:Greater than 85% as determined by SDS-PAGE.

Formulation:

The ALAD solution (1mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 100mM NaCl and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

ALAD form porphobilinogen (a precursor of heme, cytochromes and other hemoproteins) by catalyzing the compression of 2 molecules of delta-aminolevulinate. ALAD catalyzes the second step in the porphyrin and heme biosynthetic pathway; zinc is vital for enzymatic activity. ALAD has 8 identical subunits and its enzymatic activity is inhibited by lead. Mutations in the ALAD structural gene are the source for high sensitivity to lead poisoning and acute hepatic porphyria.

To place an order, please [Click HERE](#).