

HTLV-1 gp21

Description: The E.Coli derived recombinant protein contains the gp21 immunodominant region, 351-404 amino acids, the MW is 21kDa. Fused with a GST tag at N-Terminus.

Catalog #:HIPS-148

Purity: HTLV-1 gp21 is >95% pure as determined by 10% SDS-PAGE (coomassie staining) and RP-HPLC.

For research use only.

Purification Method:

HTLV-1 gp21 was purified by proprietary chromatographic technique.

Formulation:

60mM NaCl, 50mM Tris-HCl, pH-8.0, 0.25% Sarkosil, 50% Glycerol, 10mM Glutathione.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

HTLV-1 gp21 can be used as an antigen in ELISA and Western Blots. Excellent reagent for correct detection of HTLV infections, with minimal specificity problems.

Introduction:

Human T-lymphotropic virus (HTLV) is a human, single-stranded RNA retrovirus that causes T-cell leukemia and T-cell lymphoma. The virus activates a subset of T-helper cells called Th1 cells. The result is a proliferation of Th1 cells and overproduction of Th1 related cytokines (mainly IFN-gamma and TNF-alpha). Feedback mechanisms of these cytokines cause a suppression of the Th2 lymphocytes and a reduction of Th2 cytokine production (mainly IL-4, IL-5, IL-10 and IL-13). The end result is a reduction in the ability of the infected host to mount an adequate immune response to invading organisms that require a predominantly Th2 dependant response (these include parasitic infections and production of mucosal and humoral antibodies).

Storage:

HTLV-1 gp21 although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.

To place an order, please [Click HERE](#).