

HSPA5 Human

Description: Recombinant Human HSPA5 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 640 amino acids (20-650 a.a.) and having a molecular mass of 71kDa. HSPA5 human recombinant is fused to an 8 amino acid His Tag at C-terminus and purified by conventional chromatography techniques.

Catalog #: HYP5-051

For research use only.

Synonyms: BIP, MIF2, GRP78, FLJ26106, HSPA5, 78 kDa glucose-regulated protein, GRP 78, Heat shock 70 kDa protein 5, Immunoglobulin heavy chain-binding protein, Endoplasmic reticulum luminal Ca(2+)-binding protein grp78.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MEEDKKEDVG TVVGIDLGTT YSCVGVFKNG RVEIANDQG
NRITPSYVAF TPEGERLIGD AAKNQLTSNP ENTVFDKRL IGRTWNDPSV QQDIKFLPFK
VVEKTKPYI QVDIGGGQTK TFAPEEISAM VLTMMKETAE AYLGGKVTHA VVTPAYFND
AQRQATKDAG TIAGLNVMRIINEPTAAIA YGLDKREGEK NILVFDLGGG TFDVSLTID
NGVFEVVATN GDT

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

The HSPA5 protein solution contains 20mM Tris-HCl, pH-8 & 10% Glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Once Chinese hamster K12 cells are starved from glucose, the synthesis of GRP (glucose-regulated protein) expressed. HSPA5 also called BiP, is part of the HSP70 family and plays a role in the folding and assembly of proteins in the endoplasmic reticulum. HSPA5 plays a key role in monitoring protein transport through the cell. HSPA5 is a stress response protein which is induced by agents or conditions that adversely affect endoplasmic reticulum function. HSPA5 is crucial for the proper glycosylation, folding as well as for the maintenance of cell homeostasis and the prevention of apoptosis. HSPA5 is differentially expressed in the dorsolateral prefrontal cortex from patients with schizophrenia. HSPA5 guides posttranslational hepatitis B virus large envelope protein import into the mammalian ER. HSPA5 actively regulates multiple malignant phenotypes, including cell growth, migration, and invasion.

To place an order, please [Click HERE](#).