www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

SGK1 Human

Description: SGK1 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 393 amino acids (60-431 a.a.) and having a molecular mass of 44.5kDa.SGK1 is fused to a 21 amino acid His-tag at N-terminus & Durified by proprietary chromatographic techniques.

Catalog #:PKPS-279

For research use only.

Synonyms: Serine/threonine-protein kinase Sgk1, Serum/glucocorticoid-regulated kinase 1, SGK1, SGK.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MISQPQEPEL MNANPSPPPS PSQQINLGPS SNPHAKPSDF HFLKVIGKGS FGKVLLARHK AEEVFYAVKV LQKKAILKKK EEKHIMSERN VLLKNVKHPF LVGLHFSFQT ADKLYFVLDY INGGELFYHL QRERCFLEPR ARFYAAEIAS ALGYLHSLNI VYRDLKPENI LLDSQGHIVL TDFGLCKENI EHNSTTSTFC GTPEYLAPEV LH

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

SGK1 protein solution (0.25mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 30% glycerol, 0.2M NaCl, 2mM DTT and 0.1mM PMSF.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Serum-and glucocorticoid-regulated kinase (SGK1) is a serine/threonine protein kinase and a member of the AGC subfamily, which includes protein kinases A, G, and C. The SGK1 protein has an imperative role in activating specific potassium, sodium and chloride channels, suggesting a participation in the regulation of processes such as cell survival, neuronal excitability, and renal sodium excretion. SGK1 is activated in vitro by PDK-1 (3-phosphoinositide-dependent protein kinase-1) and in vivo in reaction to signals which activate phosphatidylinositol (PI) 3-kinase.

To place an order, please Click HERE.





