

## STAT1 Human

**Description:** STAT1 Recombinant Human produced in E.Coli is a single, non-glycosylated polypeptide chain containing 732 amino acids (1-712 a.a.) and having a molecular mass of 85.2 kDa. The STAT1 is fused to 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

**Catalog #:** PKPS-322

For research use only.

**Synonyms:** Signal transducer and activator of transcription 1-alpha/beta, Transcription factor ISGF-3 components p91/p84, STAT1, ISGF-3, STAT91, DKFZp686B04100.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered clear solution.

**Amino Acid Sequence:** MGSSHHHHHH SSGLVPRGSH MSQWYELQQL DSKFLEQVHQ  
LYDDSFPMEL RQYLAQWLEK QDWEHAANDV SFATIRFHD LSQLDDQYSR FSLENNFLLQ  
HNIRKSKRNL QDNFQEDPIQ MSMIYSCLK EERKILENAQ RFNQAQSGNI QSTVMLDKQK  
ELDSKVRNVK DKVMCIEHEI KSLEDLQDEY DFKCKTLQNR EHETNGVAKS DQKQEQLLLK  
KMYLMLDNKR KE

**Purity:** Greater than 90% as determined by SDS-PAGE.

**Formulation:**

STAT1 0.5mg/ml protein solution contains 20mM Tris-HCl buffer pH-8, 0.1M NaCl, 1mM DTT and 10% glycerol.

**Stability:**

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Introduction:**

STAT1 is a member of the Signal Transducers and Activators of Transcription family of transcription factors. STAT1 is involved in upregulating genes due to a signal by either type I or type II interferons. In response to IFN-stimulation, STAT1 forms homodimers or heterodimers with STAT3 that bind to the GAS (Interferon-Gamma Activated Sequence) promoter element; in response to either IFN- or IFN-stimulation, STAT1 forms a heterodimer with STAT2 that can bind the ISRE (Interferon Stimulated Response Element) promoter element. In either case, binding of the promoter element leads to an increased expression of ISG (Interferon Stimulated Genes).

**To place an order, please [Click HERE](#).**