www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

ACVR1B Human

Description: ACVR1B Human Recombinant produced in HEK-293 cells is a single, glycosylated, polypeptide chain having a molecular weight of38.2 kDa thoµgh on SDS-PAGE migrates at about 46 kDa due to the glycosilation. The ACVR1B sequence(24-126 a.a) corresponds to the extracellular region of mature Human protein, fused to the Fc region of Human IgG1.The ACVR1B is purified by proprietary chromatographic techniques.

Catalog #:PKPS-373

For research use only.

Synonyms:EC 2.7.11.30, ALK4, SKR2, ACVRLK4, ActRIB, Activin receptor type-1B.

Source: HEK293 Cells.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:SGPRGVQALL CACTSCLQAN YTCETDGACM VSIFNLDGME HHVRTCIPKV ELVPAGKPFY CLSSEDLRNT HCCYTDYCNR IDLRVPSGHL KEPEHPSMWG PVEENLYFQG GSGTKLDKTH TCPPCPAPEL LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHGDPEVK FNWYVDGVEV HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKAI PAPIFK TI

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

The ACVR1B protein was lyophilized from a 0.2

Stability:

Lyophilized ACVR1B although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution ACVR1B should be storedat 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished forLABORATORY RESEARCHUSEONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized ACVR1B in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

ACVR1B on ligand binding, creates a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. ACVR1B Phosphorylates TTRAP.

Biological Activity:

In a functional ELISA, immobilized recombinant Human ACVR1B (0.2g/mL) will bind recombinant Human ACVR1B with a linear range of 0.8 100ng/mL.

To place an order, please Click HERE.





