

SMNDC1 Human

Description: SMNDC1 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 258 amino acids (1-238a.a.) and having a molecular mass of 28.9kDa. SMNDC1 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-042

For research use only.

Synonyms: SMNR, SMN-related protein, SPF30, Survival Motor Neuron Domain Containing 1, 30kDa Splicing Factor SMNrp, Splicing Factor 30, Survival of Motor Neuron-related.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MSED LAKQLA SYKAQLQQVE
AALSGNGENE DLLKLKKDLQ EVIELTKDLL STQPSETLAS SDSFASTQPT HSWKVGDKCM
AVWSEDGQCY EAEIEEIDEE NGTAAITFAG YGNAEVTPLL NLKPVEEGRK AKEDSGNKPM
SKKEMIAQQR EYKKKKALKK AQRILEQE REDQKVKWQQ FNNRAYSKNK KGQVKRSIFA
SPESVTGKVG VG

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

The SMNDC1 protein solution (0.5mg/1ml) is formulated in 20mM Tris-HCl buffer (pH8.0) 1mM DTT, 100mM NaCl and 10% glycerol.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

SMNDC1 is a vital splicing factor obligatory for spliceosome assembly which is a part of the SMN family. SMNDC1 has one Tudor domain with significant similarity to Survival Motor Neuron (SMN) and is expressed in heart, pancreas and skeletal muscle, localizing to Cajal bodies and nuclear speckles. Mutations in SMNDC1 are cause of autosomal recessive proximal spinal muscular atrophy.

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

To place an order, please [Click HERE](#).