

## LYG2 Human

**Description:**LYG2 Human Recombinant produced in E. coli is a single polypeptide chain containing 216 amino acids (20-212) and having a molecular mass of 23.9 kDa. LYG2 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

**Catalog #:**PRPS-1252

For research use only.

**Synonyms:**Lysozyme G-Like 2, Lysozyme G-Like Protein 2, LYGH, EC 3.2.1.-.

**Source:**E.coli.

**Physical Appearance:**Sterile Filtered colorless solution.

**Amino Acid Sequence:**MGSSHHHHHH SSGLVPRGSH MGSSYPFSHS MKPHLHPRLY  
HGCGYDITM KTSGATCDAN SVMNCGIRGS EMFAEMDLRA IKPYQTLIKE VGQRHCVDPA  
VIAAIISRES HGGSVLQDGW DHRGLKFGLM QLDKQTYHPV GAWDSKEHLS QATGILTERI  
KAIQKKFPTW SVAQHLKGGL SAFKSGIEAI ATPSDIDNDF VNDIARAKF YKRQSF.

**Purity:**Greater than 90% as determined by SDS-PAGE.

### Formulation:

The LYG2 solution (0.5mg/1ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.2M NaCl, 1mM DTT and 20% glycerol.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

Lysozyme G-like 2 and LYGH (LYG2), is a member of the glycosyl hydrolase 23 family. Lysozyme takes part in human innate immunity by causing bacterial cell lysis. LYG2 contains a SLT domain, a protein domain present in bacterial lytic transglycosylase (SLT) and in eukaryotic lysozymes (GEWL). SLT domain catalyzes the cleavage of the beta-1,4-glycosidic bond between N-acetylmuramic acid (MurNAc) and N-acetylglucosamine (GlcNAc).

**To place an order, please [Click HERE](#).**