

SGCD Human

Description: SGCD Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 257 amino acids (57-289a.a) and having a molecular mass of 28kDa. SGCD is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-1380

For research use only.

Synonyms: 35DAG, CMD1L, DAGD, SG-delta, SGCDP, SGD, Delta-sarcoglycan, 35 kDa dystrophin-associated glycoprotein.

Source: E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHH SSGLVPRGSH RSHMKVMNFT IDGMGNLRIT
EKGLKLEGDS EFLQPLYAKE IQSRPGNALY FKSARNVTVN ILNDQTKVLT QLITGPKAVE
AYGKKFEVKT VSGKLLFSAD NNEVVGAER LRVLGAEGTV FPKSIETPNV RADPFKELRL
ESPTRSLVME APKGVEINAE AGNMEATCRT ELRLESKDGE IKLDAKIRL PRLPHGSYTP
TGTRQKVFEI CV

Purity: Greater than 85% as determined by SDS-PAGE.

Formulation:

SGCD protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Sarcoglycan Delta (SGCD) is one of the four known components of the sarcoglycan complex, which is a subcomplex of the dystrophin-glycoprotein complex (DGC) and is expressed mainly in skeletal and cardiac muscle. DGC forms a link between the F-actin cytoskeleton and the extracellular matrix. SGCD proteins mutations have been associated with autosomal recessive limb-girdle muscular dystrophy and dilated cardiomyopathy. SGCD gene expression is regulated by MITF in melanocytic cells.

To place an order, please [Click HERE](#).