

TIGAR Human, TAT

Description:TIGAR Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 283 amino acids (including the 270 residues of full-length TIGAR and a 13-residue C-terminal TAT peptide) and having a molecular mass of 31.7kDa. The TIGAR is purified by proprietary chromatographic techniques.

Synonyms:Fructose-2,6-bisphosphatase TIGAR, TP53-induced glycolysis and apoptosis regulator, TIGAR, C12orf5.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:MARFALTVVR HGETRFNKEK IIQQQGVDEP LSETGFKQAA
AAGIFLNNVK FTHAFSSDLM RTKQTMHGIL ERSKFCKDMT VKYDSRLRER KYGVVEGKAL
SELRAMAKAA REECPVFTPP GGETLDQVKM RGIDFFFLC QLILKEADQK EQFSQGSPTS
CLETSLAEIF PLGKNHSSKV NSDSGIPGLA ASVLVSHGA YMRSFLDYFL TDLKCSLPAT
LSRSELMSVT PN

Purity:Greater than 95.0% as determined by: (a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

TIGAR was Lyophilized from a 0.2

Stability:

Lyophilized TIGAR stable at room temperature for 3 weeks, should be stored desiccated below -18C. Upon reconstitution TIGAR should be stored at 4C between 2-7 days and for future use below -18C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized TIGAR in sterile 18M-cm H2O not less than 100

Introduction:

TIGAR is a p53-inducible enzyme which catalyzes the hydrolysis of fructose-2-6 bisphosphate (F-2-6-BP) to fructose-6-phosphate and inorganic phosphate. F-2-6-BP is an influential activator of 6-phosphofructose-1 kinase (the rate limiting enzyme of glycolysis). By lowering the intracellular level of F-2-6-BP, TIGAR expression leads to increased glucose processing through the pentose phosphate pathway, the main cellular source for NADPH. Protein transduction using TAT fusion proteins represents an alternative methodology for introducing transcription factors and other intracellular proteins into primary as well as transformed cells.

Biological Activity:

The Specific Activity was measured by its ability to protect U2OS cells from apoptosis induced by

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
hydrogen peroxide is in a concentration range of 0.1-5.0



To place an order, please [Click HERE](#).

Catalog #:PRPS-1587

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)



Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)

Request a
CATALOG
[CLICK HERE](#)