

MAPT Human

Description: MAPT Recombinant Human produced in E.Coli is a single, non-glycosylated polypeptide chain containing 372 amino acids (1-352 a.a.) and having a molecular mass of 38.9 kDa (Real molecular weight on SDS-PAGE will be shift up). The MAPT is fused to a 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

Catalog #: PRPS-302

For research use only.

Synonyms: TAU, DDPAC, FTDP-17, MAPTL, MSTD, MTBT1, MTBT2, PPND.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MAEPRQEFEV MEDHAGTYGL
GDRKDQGGYT MHQDQEGDTD AGLKAEAGI GDTPSLEDEA AGHVTQARMV SKSKDGTGSD
DKKAKGADGK TKIATPRGAA PPGQKGQANA TRIPAKTPPA PKTPSSGEP PKSGDRSGYS
SPGSPGTPGS RSRTPSLPTP PTREPKKVAV VRTPPKSPSS AKSRLQTAPV PMPDLKNVKS
KIGSTENLKH QP

Purity: Greater than 85.0% as determined by SDS-PAGE.

Formulation:

The MAPT solution (0.5mg/ml) contains 20mM Tris-HCl pH-8, 1mM DTT, 0.2M NaCl & 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

MAPT is a neuronal microtubule associated protein localized mostly on axons. MAPT promotes tubulin polymerisation and stabilizes microtubules, however it also serves to connect certain signalling pathways to the cytoskeleton. MAPT, in its hyperphosphorylated form, is the main part of paired helical filaments (PHF) and neurofibrillary lesions in Alzheimer's disease (AD) brain.

To place an order, please [Click HERE](#).