

scIC2

Description: Recombinant Single Chain Cardiac Troponin I-C 2nd generation produced in E.Coli is a single, non-glycosylated, polypeptide chain (aa28-110-Linker-cTnC), having a molecular weight of 30kDa. 1mg of scIC2 is equivalent to 0.82mg of full length human cardiac TNI on a molar basis. The scIC2 is purified by proprietary chromatographic techniques.

Catalog #: PRPS-353

For research use only.

Synonyms: Troponin I cardiac muscle, Cardiac troponin I, TNNI3, TNNC1, CMH7, RCM1, cTnI, CMD2A, MGC116817, Troponin C slow skeletal and cardiac muscles, TN-C, TNNC1, TNNC, TNC, CMD1Z.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colourless liquid formulation.

Purity: Greater than 91.0% as determined by SDS-PAGE.

Formulation:

The protein solution contains 20mM Tris-HCl, 250mM NaCl and 50% glycerol, pH 7.5.

Stability:

scIC2 although stable at 10°C for 7 days, should be stored below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

Immunoassays and western blot.

Introduction:

Troponin Complex is a heteromeric protein playing an important role in the regulation of skeletal and cardiac muscle contraction. It consists of three subunits, Troponin I, Troponin T and Troponin C. Each subunit is responsible for part of Troponin Complex function. E.g. Troponin I inhibits ATP-ase activity of acto-myosin. Troponin T and Troponin I are presented in cardiac muscles in different forms than in skeletal muscles. Purified subunits of rcTnI, rcTnC and rcTnT are recomplexed in vitro under appropriate conditions.

To place an order, please [Click HERE](#).