

## DR1 Human

**Description:** DR1 Recombinant Human produced in E.Coli is a single, non-glycosylated polypeptide chain containing 196 amino acids (1-176 a.a.) and having a molecular mass of 21.6 kDa. The DR1 is fused to 20 amino acid His-Tag at N-terminus and purified by proprietary chromatographic techniques.

**Catalog #:** PRPS-549

For research use only.

**Synonyms:** NC2, NC2-BETA, Negative cofactor 2-beta.

**Source:** Escherichia Coli.

**Amino Acid Sequence:** MGSSHHHHHH SSGLVPRGSH MASSSGNDDD LTIPRAAINK  
MIKETLPNVR VANDARELVV NCCTEFIHLI SSEANEICNK SEKKTISPEHVIQALSLGF  
GSYISEVKEV LQECKTVALK RRKASSRLN LGIPEEEELLR QQQELFAKAR QQQAELAQQE  
WLQMQQAAQQ AQLAAASASA NQAGSSQDE EDDDDI.

**Purity:** Greater than 85.0% as determined by SDS-PAGE.

**Formulation:**

DR1 protein solution (0.5mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 0.1mM PMSF, 0.1M NaCl & 10% glycerol.

**Stability:**

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Introduction:**

DR1 is a phosphoprotein that inhibits both basal and activated levels of transcription. DR1 is phosphorylated in vivo which affects its interaction with TBP. DR1 has a histone fold motif at the amino terminus, a TBP-binding domain, and a glutamine- and alanine-rich region. By selectively repressing polymerases II and III, DR1 alters the physiological balance of transcriptional output in favor of polymerase I.

**To place an order, please [Click HERE](#).**