

PAIP2 Human

Description: Recombinant Human PAIP2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 147 amino acids (1-127 a.a.) and having a molecular mass of 17.1 kDa. PAIP2 human recombinant is fused to 20 amino acid His Tag at N-terminus and purified by conventional chromatography techniques.

Catalog #: PRPS-755

For research use only.

Synonyms: Poly(A) Binding Protein Interacting Protein 2, PABP- interacting protein 2, HSPC218, PAIP-2, PAIP2A, Polyadenylate-binding protein-interacting protein 2, PAIP2, MGC72018.

Source: Escherichia Coli.

Physical Appearance: Sterile filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MKDPSRSSTS PSINEDVII
NGSHEDDNP FAEYMWME NE EEFNRQIEEE LWEEEFIERC FQEMLEEEEE HEWFIPARDL
PQTMDIQDQ FNDLVISDGS SLEDLVVKS NLPNAKEFVP GVKYGN I.

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

The PAIP2 protein solution contains 20mM Tris-HCl, pH-8, 1mM DTT & 10% Glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

PAIP2 has a role as a repressor in the regulation of translation initiation of poly(A)-containing mRNAs. PAIP2 inhibitory activity on translation is mediated through its action on PABPC1. PAIP2 displaces the interaction of PABPC1 with poly(A) RNA and competes with PAIP1 for binding to PABPC1. PAIP2 association with PABPC1 results in disruption of the cytoplasmic poly(A) RNP structure organization.

To place an order, please [Click HERE](#).