

MYCBP Human

Description: MYCBP Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 123 amino acids (1-103 a.a.) and having a molecular mass of 14.1kDa. MYCBP is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-949

For research use only.

Synonyms: C-Myc-binding protein, Associate of Myc 1, AMY-1, MYCBP, AMY1, FLJ41056.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MAHYKAADSK REQFRYLEK
SGVLDTLTKV LVALYEEPEK PNSALDFLKH HLGAATPENP EIELLRLELA EMKEKYEIV
EENKCLKAKL AQYEPPQEEK RAE.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

MYCBP protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 20% glycerol, 0.1M NaCl and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

MYCBP is a member of the AMY1 family. MYCBP (c-Myc binding protein) binds to the transactivation domain of c-Myc and stimulates the activation of E-box-dependent transcription. MYCBP translocates from the cytoplasm to the nucleus during S phase when increased expression of c-Myc occurs. MYCBP also associates with AKAP 149 and AKAP 84 in mitochondria of somatic cells and sperm, suggesting a role for MYCBP in spermatogenesis. MYCBP is highly expressed in the heart, placenta, pancreas, skeletal muscle and kidney. It is also present at low levels in the lung.

To place an order, please [Click HERE](#).