www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

GP2 Human

Description: GP2 is a cDNA coding for the free form of the human pancreatic secretory granule membrane major glycoprotein / GP2 having a molecular mass of 54,909 Dalton (protein component excluding glycosylation; observed molecular weight approx. 68 kDa) pH 5.4. GP2 protein is fused to a deca-histidine purification tag.

Catalog #:prPS-122

For research use only.

Synonyms: Glycoprotein 2 (zymogen granule membrane), Pancreatic zymogen granule membrane protein GP-2, ZAP75, DKFZp779K0533.

Source:Sf9 insect cells.

Purity: Greater than 80% as determined by SDS-PAGE.

Formulation:

GP2 is supplied in 16mM HEPES buffer pH-8.0, 400mM NaCl, and 20% glycerol.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

Western blot with monoclonal anti-GP2 antibodies and monoclonal anti-hexa-His-tag antibody.

Introduction:

Crohn's Disease (CD), which is very common in Caucasians, is an inflammatory bowel diseases (IBD). Mucosal inflammation in CD seems to arise when dysregulation of the immune system breaks the balance between tolerance to commensal microbiota or food-derived antigens and immunity to pathogens. The autoimmune mechanisms takes part in the development of CD, and exocrine pancreas autoantibodies (PAB) are disease-specific in CD patients. It was established not too long ago that glycoprotein 2 (GP2) is the major autoantigenic mark recognized by CD-specific PAB. Furthermore to IgG and IgM PAB isotypes, IgA pancreatic autoantibodies were also identified in CD patients. GP2 is an extremely glycosylated 78 kDa protein with N-linked carbohydrates. It is responsible for up to 40% of all zymogen granule (ZG) membrane proteins in pancreatic acinar cells and is linked to the ZG membrane via a glycosyl phosphoinositol (GPI) anchor.

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

To place an order, please Click HERE.





